

DFUB 98-5
 TP-USL/98/4
 May 1998

The Master Differential Equations for the 2-loop Sunrise Selfmass Amplitudes.

M. Caffo^{ab}, H. Czyż^{c*}, S. Laporta^b and E. Remiddi^{ba}

^a INFN, Sezione di Bologna, I-40126 Bologna, Italy

^b Dipartimento di Fisica, Università di Bologna, I-40126 Bologna, Italy

^c Institute of Physics, University of Silesia, PL-40007 Katowice, Poland

e-mail: caffo@bo.infn.it

czyz@usctoux1.cto.us.edu.pl

laporta@bo.infn.it

remiddi@bo.infn.it

Abstract

The master differential equations in the external square momentum p^2 for the master integrals of the two-loop sunrise graph, in n -continuous dimensions and for arbitrary values of the internal masses, are derived. The equations are then used for working out the values at $p^2 = 0$ and the expansions in p^2 at $p^2 = 0$, in $(n - 4)$ at $n \rightarrow 4$ limit and in $1/p^2$ for large values of p^2 .

PACS 11.10.-z Field theory

PACS 11.10.Kk Field theories in dimensions other than four

PACS 11.15.Bt General properties of perturbation theory

*Partly supported by the Polish Committee for Scientific Research under grant no 2P03B17708.

1 Introduction.

In a previous paper [1], it was shown that a linear system of first order differential equations in the external Mandelstam variables, to be referred to from now on as master equations, can in general be written for the master integrals of any Feynman graph. In this paper the master equations are written for the 2-loop sunrise self-mass graph amplitudes in n continuous dimensions and for arbitrary values of the masses as a function of the square momentum transfer p^2 . All momenta are Euclidean, so that $p^2 > 0$ corresponds to spacelike p_μ . The master equations can be used for obtaining valuable information on the properties of the master integrals, for working out virtually any desired expansion, for obtaining closed analytic results (when possible) and, in any case, for the actual numerical calculation of the amplitudes.

In the various sections of this paper we will show

- how to use the integration by part identities [2] , [3] for writing the master equations;
- how to obtain the (analytic) values of the master integrals at $p^2 = 0$, in the $n \rightarrow 4$ limit;
- how to expand the master integrals in p^2 at $p^2 = 0$;
- how to expand the master integrals in $(n - 4)$ for arbitrary p^2 ;
- how to solve analytically the master equations when two masses are equal to zero;
- how to expand the master integrals in $1/p^2$ for large values of p^2 and arbitrary values of the masses.

The appendix contains a few ready-to-use recurrence formulae, obtained by solving the integration by parts identities for the considered graph [3].

All the algebra needed to generate the identities, the equations and the considered expansions has been processed by means of the computer program FORM [4] by J. Vermaseren.

While we deal with the 2 loop case only, it should be apparent from the discussion that some suitable extension of the results may apply also to more general cases.

2 The master integrals and their differential equations.

The 2-loop sunrise self-mass graph is shown in Fig.1.




Figure 1: The 2-loop sunrise self-mass graph.

The related amplitudes, in n continuous dimensions and fully Euclidean variables, can be written as

$$A(n, m_1^2, m_2^2, m_3^2, p^2, -\alpha_1, -\alpha_2, -\alpha_3, \beta_1, \beta_2) = \int \frac{d^n k_1}{(2\pi)^{n-2}} \int \frac{d^n k_2}{(2\pi)^{n-2}} \frac{(p \cdot k_1)^{\beta_1} (p \cdot k_2)^{\beta_2}}{(k_1^2 + m_1^2)^{\alpha_1} (k_2^2 + m_2^2)^{\alpha_2} ((p - k_1 - k_2)^2 + m_3^2)^{\alpha_3}} . \quad (1)$$

It is by now well known that, as a consequence of the integration by parts identities [2], all the above amplitudes can be expressed in terms of four master integrals [3], which can be taken as

$$\begin{aligned} F_0(n, m_1^2, m_2^2, m_3^2, p^2) &= A(n, m_1^2, m_2^2, m_3^2, p^2, -1, -1, -1, 0, 0) , \\ F_1(n, m_1^2, m_2^2, m_3^2, p^2) &= A(n, m_1^2, m_2^2, m_3^2, p^2, -2, -1, -1, 0, 0) , \\ F_2(n, m_1^2, m_2^2, m_3^2, p^2) &= A(n, m_1^2, m_2^2, m_3^2, p^2, -1, -2, -1, 0, 0) , \\ F_3(n, m_1^2, m_2^2, m_3^2, p^2) &= A(n, m_1^2, m_2^2, m_3^2, p^2, -1, -1, -2, 0, 0) . \end{aligned} \quad (2)$$

It follows from the definitions that the last three master integrals are the mass derivatives of the first, i.e. for $i = 1, 2, 3$

$$F_i(n, m_1^2, m_2^2, m_3^2, p^2) = -\frac{\partial}{\partial m_i^2} F_0(n, m_1^2, m_2^2, m_3^2, p^2) . \quad (3)$$

The simplest way for deriving the required differential equations for self-mass amplitudes, which depend on a single Mandelstam variable only, consists in writing for them the familiar scaling equation (which can be obtained, for instance, by scaling in the definition Eq.(1) all the integration momenta by a same factor λ and then differentiating with respect to λ).

For $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ that procedure gives

$$\left(p^2 \frac{\partial}{\partial p^2} + m_1^2 \frac{\partial}{\partial m_1^2} + m_2^2 \frac{\partial}{\partial m_2^2} + m_3^2 \frac{\partial}{\partial m_3^2} - (n-3) \right) F_0(n, m_1^2, m_2^2, m_3^2, p^2) = 0, \quad (4)$$

i.e., on account of Eq.(3),

$$\begin{aligned} p^2 \frac{\partial}{\partial p^2} F_0(n, m_1^2, m_2^2, m_3^2, p^2) &= (n-3) F_0(n, m_1^2, m_2^2, m_3^2, p^2) + m_1^2 F_1(n, m_1^2, m_2^2, m_3^2, p^2) \\ &+ m_2^2 F_2(n, m_1^2, m_2^2, m_3^2, p^2) + m_3^2 F_3(n, m_1^2, m_2^2, m_3^2, p^2). \end{aligned} \quad (5)$$

A similar equation holds for each of $F_i(n, m_1^2, m_2^2, m_3^2, p^2)$, $i = 1, 2, 3$

$$\left(p^2 \frac{\partial}{\partial p^2} + m_1^2 \frac{\partial}{\partial m_1^2} + m_2^2 \frac{\partial}{\partial m_2^2} + m_3^2 \frac{\partial}{\partial m_3^2} - (n-4) \right) F_i(n, m_1^2, m_2^2, m_3^2, p^2) = 0. \quad (6)$$

When carrying out the mass derivatives, new integrals appear, like for instance those corresponding to $\alpha_1 = 3$, $\alpha_2 = \alpha_3 = 1$ and $\beta_1 = \beta_2 = 0$ in the notation of Eq.(1). All those integrals, according to the formulas of the Appendix, can be expressed in terms of the master integrals. The result can be written as

$$\begin{aligned} p^2 \frac{\partial}{\partial p^2} F_i(n, m_1^2, m_2^2, m_3^2, p^2) &= \frac{n-4}{2} F_i(n, m_1^2, m_2^2, m_3^2, p^2) \\ &+ \frac{n-3}{D(m_1^2, m_2^2, m_3^2, p^2)} \left[\begin{aligned} &P_{i,0}(m_1^2, m_2^2, m_3^2, p^2) \frac{3n-8}{2} F_0(n, m_1^2, m_2^2, m_3^2, p^2) \\ &+ P_{i,1}(m_1^2, m_2^2, m_3^2, p^2) F_1(n, m_1^2, m_2^2, m_3^2, p^2) \\ &+ P_{i,2}(m_1^2, m_2^2, m_3^2, p^2) F_2(n, m_1^2, m_2^2, m_3^2, p^2) \\ &+ P_{i,3}(m_1^2, m_2^2, m_3^2, p^2) F_3(n, m_1^2, m_2^2, m_3^2, p^2) \end{aligned} \right] \\ &+ \frac{(n-2)^2}{D(m_1^2, m_2^2, m_3^2, p^2)} \left[\begin{aligned} &Q_{i,1}(m_1^2, m_2^2, m_3^2, p^2) T(n, m_2^2) T(n, m_3^2) \\ &+ Q_{i,2}(m_1^2, m_2^2, m_3^2, p^2) T(n, m_1^2) T(n, m_3^2) \\ &+ Q_{i,3}(m_1^2, m_2^2, m_3^2, p^2) T(n, m_1^2) T(n, m_2^2) \end{aligned} \right], \end{aligned} \quad (7)$$

where $D(m_1^2, m_2^2, m_3^2, p^2)$, the $P_{i,j}(m_1^2, m_2^2, m_3^2, p^2)$ and the $Q_{i,j}(m_1^2, m_2^2, m_3^2, p^2)$ are polynomials in m_1^2, m_2^2, m_3^2 and p^2 , to be discussed in a moment, while $T(n, m^2)$ is defined by

$$T(n, m^2) = \int \frac{d^n k}{(2\pi)^{n-2}} \frac{1}{k^2 + m^2} = \frac{m^{n-2}}{(n-2)(n-4)} C(n) , \quad (8)$$

$C(n)$ being the following function of n

$$C(n) = \left(2\sqrt{\pi}\right)^{(4-n)} \Gamma\left(3 - \frac{n}{2}\right) . \quad (9)$$

The expansion of Eq.(9) in n around $n = 4$ is

$$\begin{aligned} C(n) &= 1 - (n-4) \left[\log(2\sqrt{\pi}) - \frac{1}{2}\gamma_E \right] \\ &+ (n-4)^2 \frac{1}{2} \left[\log^2(2\sqrt{\pi}) + \frac{1}{4} \left(\frac{\pi^2}{6} + \gamma_E^2 \right) - \gamma_E \log(2\sqrt{\pi}) \right] \\ &+ \mathcal{O}\left((n-4)^3\right) , \end{aligned} \quad (10)$$

where γ_E is the Euler-Mascheroni constant. The expansion, which is given only to make easier the comparisons with other results in the literature, is however never needed in practice, despite the frequent presence of $1/(n-4)$ factors, as in Eq.(8). Indeed, there is always an overall factor $C(n)$ for each loop; in the calculation of any physical (and therefore finite) quantity, the singularities, which are of the form $C^2(n)/(n-4)^2$ etc., cancel out, and in the remaining finite part it is sufficient to use the value at $n = 4$, which is simply

$$C(4) = 1 . \quad (11)$$

The explicit expression of the polynomial $D(m_1^2, m_2^2, m_3^2, p^2)$ is

$$\begin{aligned} D(m_1^2, m_2^2, m_3^2, p^2) &= (p^2)^4 + 4(m_1^2 + m_2^2 + m_3^2)(p^2)^3 \\ &+ 2\left(3m_1^4 + 3m_2^4 + 3m_3^4 + 2m_1^2 m_2^2 + 2m_1^2 m_3^2 + 2m_2^2 m_3^2\right)(p^2)^2 \\ &+ 4(m_1^6 + m_2^6 + m_3^6 - m_1^4 m_2^2 - m_1^4 m_3^2 - m_1^2 m_2^4 - m_1^2 m_3^4 - m_2^4 m_3^2 - m_2^2 m_3^4 \\ &\quad + 10m_1^2 m_2^2 m_3^2) p^2 + R^4(m_1^2, m_2^2, m_3^2) \\ &= \left(p^2 + (m_1 + m_2 + m_3)^2\right) \left(p^2 + (m_1 + m_2 - m_3)^2\right) \\ &\quad \cdot \left(p^2 + (m_1 - m_2 + m_3)^2\right) \left(p^2 + (m_1 - m_2 - m_3)^2\right) , \end{aligned} \quad (12)$$

where

$$\begin{aligned} R^2(m_1^2, m_2^2, m_3^2) &= m_1^4 + m_2^4 + m_3^4 - 2m_1^2 m_2^2 - 2m_1^2 m_3^2 - 2m_2^2 m_3^2 \\ &= (m_1 + m_2 + m_3)(m_1 + m_2 - m_3) \\ &\quad \cdot (m_1 - m_2 + m_3)(m_1 - m_2 - m_3) . \end{aligned} \quad (13)$$

The polynomial $P_{i,j}$, for $i = 1$ are given by

$$\begin{aligned}
P_{1,0}(m_1^2, m_2^2, m_3^2, p^2) &= -(p^2)^3 - (3m_1^2 + m_2^2 + m_3^2)(p^2)^2 \\
&\quad - (3m_1^4 - m_2^4 - m_3^4 - 2m_1^2m_2^2 - 2m_1^2m_3^2 + 10m_2^2m_3^2)p^2 \\
&\quad - (m_1^2 - m_2^2 - m_3^2)R^2(m_1^2, m_2^2, m_3^2) , \\
P_{2,0}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,0}(m_2^2, m_1^2, m_3^2, p^2) , \\
P_{3,0}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,0}(m_3^2, m_2^2, m_1^2, p^2) ,
\end{aligned} \tag{14}$$

$$\begin{aligned}
P_{1,1}(m_1^2, m_2^2, m_3^2, p^2) &= -(2m_1^2 + m_2^2 + m_3^2)(p^2)^3 \\
&\quad - (6m_1^4 + 3m_2^4 + 3m_3^4 + 3m_1^2m_2^2 + 3m_1^2m_3^2 + 2m_2^2m_3^2)(p^2)^2 \\
&\quad - (6m_1^6 + 3m_2^6 + 3m_3^6 - 5m_1^4m_2^2 - 5m_1^4m_3^2 - 4m_1^2m_2^4 - 4m_1^2m_3^4 \\
&\quad - 3m_2^4m_3^2 - 3m_2^2m_3^4 + 40m_1^2m_2^2m_3^2)p^2 \\
&\quad - (2m_1^4 + m_2^4 + m_3^4 - 3m_1^2m_2^2 - 3m_1^2m_3^2 - 2m_2^2m_3^2)R^2(m_1^2, m_2^2, m_3^2) , \\
P_{2,2}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,1}(m_2^2, m_1^2, m_3^2, p^2) , \\
P_{3,3}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,1}(m_3^2, m_2^2, m_1^2, p^2) ,
\end{aligned} \tag{15}$$

$$\begin{aligned}
P_{1,2}(m_1^2, m_2^2, m_3^2, p^2) &= m_2^2 \left[-3(p^2)^3 - (7m_1^2 + 5m_2^2 - 3m_3^2)(p^2)^2 \right. \\
&\quad - (5m_1^4 + m_2^4 - 7m_3^4 - 6m_1^2m_2^2 + 2m_1^2m_3^2 + 14m_2^2m_3^2)p^2 \\
&\quad \left. - (m_1^2 - m_2^2 - m_3^2)R^2(m_1^2, m_2^2, m_3^2) \right] ,
\end{aligned} \tag{16}$$

$$\begin{aligned}
P_{1,3}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,2}(m_1^2, m_3^2, m_2^2, p^2) , \\
P_{2,1}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,2}(m_2^2, m_1^2, m_3^2, p^2) , \\
P_{2,3}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,2}(m_2^2, m_3^2, m_1^2, p^2) , \\
P_{3,1}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,2}(m_3^2, m_1^2, m_2^2, p^2) , \\
P_{3,2}(m_1^2, m_2^2, m_3^2, p^2) &= P_{1,2}(m_3^2, m_2^2, m_1^2, p^2) ;
\end{aligned} \tag{17}$$

similarly, the $Q_{i,j}$ are given by

$$\begin{aligned}
Q_{1,1}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{2,2}(m_1^2, m_2^2, m_3^2, p^2) = Q_{3,3}(m_1^2, m_2^2, m_3^2, p^2) = \\
&= -\frac{1}{2} \left[3(p^2)^2 + 2(m_1^2 + m_2^2 + m_3^2)p^2 - R^2(m_1^2, m_2^2, m_3^2) \right] ,
\end{aligned} \tag{18}$$

$$Q_{1,2}(m_1^2, m_2^2, m_3^2, p^2) = \frac{1}{4m_1^2} \left[(p^2)^3 + (m_1^2 + m_2^2 + 3m_3^2)(p^2)^2 \right.$$

$$\begin{aligned}
& - (m_1^4 + m_2^4 - 3m_3^4 - 10m_1^2m_2^2 + 2m_1^2m_3^2 + 2m_2^2m_3^2)p^2 \\
& - (m_1^2 + m_2^2 - m_3^2)R^2(m_1^2, m_2^2, m_3^2) \Big] , \tag{19}
\end{aligned}$$

$$\begin{aligned}
Q_{1,3}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{1,2}(m_1^2, m_3^2, m_2^2, p^2) , \\
Q_{2,1}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{1,2}(m_2^2, m_1^2, m_3^2, p^2) , \\
Q_{2,3}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{1,2}(m_2^2, m_3^2, m_1^2, p^2) , \\
Q_{3,1}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{1,2}(m_3^2, m_1^2, m_2^2, p^2) , \\
Q_{3,2}(m_1^2, m_2^2, m_3^2, p^2) &= Q_{1,2}(m_3^2, m_2^2, m_1^2, p^2) . \tag{20}
\end{aligned}$$

The values corresponding to the other indices can be obtained by a suitable permutation of the masses.

The above polynomials satisfy the relations

$$\begin{aligned}
m_2^2m_3^2 Q_{1,1}(m_1^2, m_2^2, m_3^2, p^2) &= -\frac{1}{4}(m_1^2 - m_2^2 - m_3^2)P_{1,0}(m_1^2, m_2^2, m_3^2, p^2) \\
&+ \frac{1}{8} \left(P_{1,1}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,2}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,3}(m_1^2, m_2^2, m_3^2, p^2) \right) , \\
m_1^2m_3^2 Q_{1,2}(m_1^2, m_2^2, m_3^2, p^2) &= \frac{1}{4}(m_1^2 - m_2^2 + m_3^2)P_{1,0}(m_1^2, m_2^2, m_3^2, p^2) \\
&- \frac{1}{8} \left(P_{1,1}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,2}(m_1^2, m_2^2, m_3^2, p^2) + P_{1,3}(m_1^2, m_2^2, m_3^2, p^2) \right) , \\
m_1^2m_2^2 Q_{1,3}(m_1^2, m_2^2, m_3^2, p^2) &= \frac{1}{4}(m_1^2 + m_2^2 - m_3^2)P_{1,0}(m_1^2, m_2^2, m_3^2, p^2) \\
&- \frac{1}{8} \left(P_{1,1}(m_1^2, m_2^2, m_3^2, p^2) + P_{1,2}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,3}(m_1^2, m_2^2, m_3^2, p^2) \right) , \tag{21}
\end{aligned}$$

and

$$\begin{aligned}
D(m_1^2, m_2^2, m_3^2, p^2) &= -(p^2 - m_1^2 - m_2^2 - m_3^2)P_{1,0}(m_1^2, m_2^2, m_3^2, p^2) \\
&- P_{1,1}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,2}(m_1^2, m_2^2, m_3^2, p^2) - P_{1,3}(m_1^2, m_2^2, m_3^2, p^2) . \tag{22}
\end{aligned}$$

The equations Eq.s(5,7) are the master differential equations referred to in the title of this paper. They are long, but their structure is simple: they express the derivatives with respect to p^2 of the four master integrals as a linear combination of the master integrals themselves and of known non-homogenous terms, the coefficients being ratios of known polynomials in the masses and p^2 . The known terms are just products of the 1-loop, 1-denominator vacuum amplitude $T(n, m^2)$ of Eq.(8), the simplest integral in the game which fixes the overall normalization.

All the coefficients of the linear combinations are equal to a simple factor, depending on the dimension n times a ratio of polynomials in p^2 and the masses. There is a unique denominator, the polynomial $D(m_1^2, m_2^2, m_3^2, p^2)$ of Eq.(12); note that its zeros in p^2 correspond to the threshold and the three pseudo-thresholds of the considered 2-loop sunrise graph.

It should be apparent, from the derivation, that master equations of similar structure, (*i.e.* equations whose coefficients are ratios of polynomials) can be established for the master integrals of virtually any Feynman graph, for any number of loops; the size of the polynomials is expected to grow quickly with the number of the denominators and of the loops, but they are in any case finite polynomials, which can be conveniently dealt with by means of an algebraic program. Quite in general, the non-homogeneous terms correspond to amplitudes of related Feynman graphs in which one of the propagators is missing; from this point of view, the l -loop sunrise amplitudes are the simplest l -loop amplitudes, in the sense that in their equations the non-homogeneous terms are just products of $l - 1$ 1-denominator vacuum amplitudes, Eq.(8). Sunrise amplitudes are in turn present as non-homogeneous terms in the equations for the amplitudes of graphs having one more propagator and so on.

As it will be shown in the rest of the paper, the master equations contain virtually all the information needed for studying the analytic properties of the master integrals and for obtaining any expansion of interest; furthermore, after some minor work (the expansion in $(n-4)$ at $n = 4$), the differential equations can be used for the actual numerical evaluation of the master integrals, without explicit reference to the original integral representation Eq.(1), once the values of the integrals are known at some initial value of p^2 . Both the expansion in $(n-4)$ and the evaluation of the initial values (at $p^2 = 0$), can be carried out by exploiting the information contained in the master equations (which hold identically in n and are regular at $p^2 = 0$ despite the appearance of an apparent kinematical singularity at that value of p^2).

3 The value at $p^2 = 0$.

From the definitions Eq.s(1,2), it is obvious that all the master integrals as well as all their derivatives are regular at $p^2 = 0$ (which in the considered Euclidean region corresponds to $p_\mu = 0$). If we introduce the vacuum graph of Fig.2




Figure 2: The 2-loop sunrise vacuum graph.

and the corresponding amplitude

$$V(n, m_1^2, m_2^2, m_3^2) = \int \frac{d^n k_1}{(2\pi)^{n-2}} \int \frac{d^n k_2}{(2\pi)^{n-2}} \frac{1}{(k_1^2 + m_1^2)(k_2^2 + m_2^2)((k_1 + k_2)^2 + m_3^2)} , \quad (23)$$

the $p^2 = 0$ values of the master integrals defined in Eq.s(2) can then be written as

$$F_0(n, m_1^2, m_2^2, m_3^2, 0) = V(n, m_1^2, m_2^2, m_3^2) , \quad (24)$$

$$F_i(n, m_1^2, m_2^2, m_3^2, 0) = -\frac{\partial}{\partial m_i^2} V(n, m_1^2, m_2^2, m_3^2) . \quad (25)$$

Let us now consider the differential equations Eq.(5) and Eq.(7) for, say, $i = 3$ at $p^2 = 0$. Due to the regularity of the derivatives, the left hand sides of both equations vanish; therefore, the right hand sides must also vanish at $p^2 = 0$. It turns out that both equations involve the same combination of $F_1(n, m_1^2, m_2^2, m_3^2, 0)$ and $F_2(n, m_1^2, m_2^2, m_3^2, 0)$, namely $m_1^2 F_1(n, m_1^2, m_2^2, m_3^2, 0) + m_2^2 F_2(n, m_1^2, m_2^2, m_3^2, 0)$; by eliminating that combination, one is left with a single equation involving only $F_0(n, m_1^2, m_2^2, m_3^2, 0)$ and $F_3(n, m_1^2, m_2^2, m_3^2, 0)$, which is its derivative with respect to m_3^2 , according to Eq.(25) ; in terms of the above vacuum amplitude the equation reads

$$\begin{aligned} R^2(m_1^2, m_2^2, m_3^2) \frac{\partial}{\partial m_3^2} V(n, m_1^2, m_2^2, m_3^2) = & \\ & (n-3)(m_3^2 - m_1^2 - m_2^2)V(n, m_1^2, m_2^2, m_3^2) \\ & + (n-2) \frac{m_3^2 - m_1^2 + m_2^2}{2m_3^2} T(n, m_1^2) T(n, m_2^2) \\ & + (n-2) \frac{m_3^2 + m_1^2 - m_2^2}{2m_3^2} T(n, m_2^2) T(n, m_3^2) \\ & - (n-2) T(n, m_1^2) T(n, m_2^2) , \end{aligned} \quad (26)$$

where $R^2(m_1^2, m_2^2, m_3^2)$ is the same as in Eq.(13). Eq.(26) is the master equation for the 2-loop vacuum amplitude of Fig.(2) – this time in one of the masses (the differential equations in the other masses are immediately obtained by a suitable permutation of the masses themselves). It is also easy to see that the same equations can be obtained by writing down directly the integration by part identities for $V(n, m_1^2, m_2^2, m_3^2)$, Eq.(23) and the related amplitudes.

Eq.(26) can be used for evaluating $V(n, m_1^2, m_2^2, m_3^2)$ once its “initial” value corresponding to a particular value of m_3^2 is given. Note that the equation itself can provide such a value. Indeed, the vacuum amplitude and all its derivatives are regular for non-vanishing values of the masses; by considering the particular value $m_3 = m_1 + m_2$, the factor $R^2(m_1^2, m_2^2, m_3^2)$ on the *l.h.s.* of Eq.(26) vanish, so that the *r.h.s.* must also vanish; that amounts to the relation

$$V(n, m_1^2, m_2^2, (m_1 + m_2)^2) = \frac{n-2}{2(n-3)} \left[\frac{T(n, m_1^2)}{m_1} \frac{T(n, m_2^2)}{m_2} - \left(\frac{T(n, m_1^2)}{m_1} + \frac{T(n, m_2^2)}{m_2} \right) \frac{T(n, (m_1 + m_2)^2)}{(m_1 + m_2)} \right]. \quad (27)$$

Eq.(27) can be taken as the “initial condition” which determines completely the solution of Eq.(26). Therefore, the differential equation Eq.(26), a first order equation in a single unknown function, can be solved in quadrature using the initial value given by Eq.(27).

For the explicit calculations it is convenient to expand the amplitude $V(n, m_1^2, m_2^2, m_3^2)$ around $n = 4$, where it is expected to develop polar singularities in $(n - 4)$. More exactly, Eq.(8) shows that $T(n, m_i^2)$ has a single pole $(n - 4)$, so that the non homogenous part of Eq.(26) and the initial value $V(n, m_1^2, m_2^2, (m_1 + m_2)^2)$, Eq.(27), both develop double poles; it follows that also $V(n, m_1^2, m_2^2, m_3^2)$ can have at most double poles. It can therefore be expanded as

$$V(n, m_1^2, m_2^2, m_3^2) = C^2(n) \left\{ \frac{1}{(n-4)^2} V^{(-2)}(m_1^2, m_2^2, m_3^2) + \frac{1}{(n-4)} V^{(-1)}(m_1^2, m_2^2, m_3^2) + V^{(0)}(m_1^2, m_2^2, m_3^2) + \mathcal{O}(n-4) \right\}, \quad (28)$$

where $C(n)$ is given by Eq.(8) and $C^2(n)$ has been treated as an overall factor according to the remarks preceding Eq.(11). Correspondingly, Eq.(26) splits into the following set of equations for the residua and the finite part at $n = 4$

$$R^2(m_1^2, m_2^2, m_3^2) - \frac{\partial}{\partial m_3^2} V^{(-2)}(m_1^2, m_2^2, m_3^2) = (m_3^2 - m_1^2 - m_2^2) V^{(-2)}(m_1^2, m_2^2, m_3^2) - \frac{1}{4} [m_1^2(m_1^2 - m_3^2) + m_2^2(m_2^2 - m_3^2)], \quad (29)$$

$$R^2(m_1^2, m_2^2, m_3^2) - \frac{\partial}{\partial m_3^2} V^{(-1)}(m_1^2, m_2^2, m_3^2) = (m_3^2 - m_1^2 - m_2^2) V^{(-1)}(m_1^2, m_2^2, m_3^2) + (m_3^2 - m_1^2 - m_2^2) V^{(-2)}(m_1^2, m_2^2, m_3^2)$$

$$\begin{aligned}
& + \frac{1}{8} \left\{ m_1^2(m_1^2 - m_3^2) + m_2^2(m_2^2 - m_3^2) \right. \\
& - \left[m_1^2 \log(m_1^2) + m_2^2 \log(m_2^2) \right] (m_1^2 + m_2^2 - m_3^2) \\
& \left. - \log(m_3^2) \left[m_1^2(m_1^2 - m_2^2 - m_3^2) + m_2^2(-m_1^2 + m_2^2 - m_3^2) \right] \right\} , \quad (30)
\end{aligned}$$

$$\begin{aligned}
R^2(m_1^2, m_2^2, m_3^2) & - \frac{\partial}{\partial m_3^2} V^{(0)}(m_1^2, m_2^2, m_3^2) = (m_3^2 - m_1^2 - m_2^2) V^{(0)}(m_1^2, m_2^2, m_3^2) \\
& + (m_3^2 - m_1^2 - m_2^2) V^{(-1)}(m_1^2, m_2^2, m_3^2) \\
& + \frac{1}{16} \left\{ -m_1^2(m_1^2 - m_3^2) - m_2^2(m_2^2 - m_3^2) - 2m_1^2 m_2^2 \log(m_1^2) \log(m_2^2) \right. \\
& + \left[m_1^2 \log(m_1^2) \left(1 - \frac{1}{2} \log(m_1^2) \right) + m_2^2 \log(m_2^2) \left(1 - \frac{1}{2} \log(m_2^2) \right) \right] (m_1^2 + m_2^2 - m_3^2) \\
& + \log(m_3^2) \left(1 - \frac{1}{2} \log(m_3^2) \right) \left[m_1^2(m_1^2 - m_2^2 - m_3^2) + m_2^2(-m_1^2 + m_2^2 - m_3^2) \right] \\
& \left. - \log(m_3^2) \left[m_1^2(m_1^2 - m_2^2 - m_3^2) \log(m_1^2) + m_2^2(-m_1^2 + m_2^2 - m_3^2) \log(m_2^2) \right] \right\} . \quad (31)
\end{aligned}$$

Similarly, in the notation of Eq.(28) the coefficients of the expansion of Eq.(27) are

$$\begin{aligned}
V^{(-2)}(m_1^2, m_2^2, (m_1 + m_2)^2) & = -\frac{1}{4}(m_1 m_2 + m_1^2 + m_2^2) , \\
V^{(-1)}(m_1^2, m_2^2, (m_1 + m_2)^2) & = \frac{3}{8}(m_1 m_2 + m_1^2 + m_2^2) \\
& - \frac{1}{4} \left[m_1^2 \log(m_1) + m_2^2 \log(m_2) + (m_1 + m_2)^2 \log(m_1 + m_2) \right] , \\
V^{(0)}(m_1^2, m_2^2, (m_1 + m_2)^2) & = -\frac{7}{16}(m_1 m_2 + m_1^2 + m_2^2) \\
& + \frac{3}{8} \left[m_1^2 \log(m_1) + m_2^2 \log(m_2) + (m_1 + m_2)^2 \log(m_1 + m_2) \right] \\
& - \frac{1}{8} [m_1 \log(m_1) - m_2 \log(m_2)]^2 - \frac{1}{8} (m_1 + m_2)^2 \log^2(m_1 + m_2) \\
& - \frac{1}{4} [m_1 \log(m_1) + m_2 \log(m_2)] (m_1 + m_2) \log(m_1 + m_2) . \quad (32)
\end{aligned}$$

When solved recursively, starting from $j = -2$, the equations (29,30,31) are all of the form

$$\begin{aligned}
R^2(m_1^2, m_2^2, m_3^2) \frac{\partial}{\partial m_3^2} V^{(j)}(m_1^2, m_2^2, m_3^2) & = (m_3^2 - m_1^2 - m_2^2) V^{(j)}(m_1^2, m_2^2, m_3^2) \\
& + g^{(j)}(m_1^2, m_2^2, m_3^2) , \quad (33)
\end{aligned}$$

where the $g^{(j)}(m_1^2, m_2^2, m_3^2)$, $j = -2, -1, 0$, to be identified by comparison with the original equations (29,30,31), can be considered as known.

Eq.(33) can in principle be solved by the quadrature formula

$$\begin{aligned} V^{(j)}(m_1^2, m_2^2, m_3^2) &= R(m_1^2, m_2^2, m_3^2) \left\{ \frac{V^{(j)}(m_1^2, m_2^2, z)}{R(m_1^2, m_2^2, z)} \right. \\ &\quad \left. + \int_z^{m_3^2} \frac{dx}{R^3(m_1^2, m_2^2, x)} g^{(j)}(m_1^2, m_2^2, x) \right\}, \end{aligned} \quad (34)$$

where z is any initial point. Our initial point is $(m_1 + m_2)^2$; as for that value of z the two terms in the *r.h.s.* diverge, it is convenient to modify the formula by carrying out an integration by parts and then to take the $z = (m_1 + m_2)^2$ limit. When that is done Eq.(34) becomes

$$\begin{aligned} V^{(j)}(m_1^2, m_2^2, m_3^2) &= \frac{1}{4m_1^2 m_2^2} \left\{ R(m_1^2, m_2^2, m_3^2) \int_{(m_1+m_2)^2}^{m_3^2} dx \frac{x - m_1^2 - m_2^2}{R(m_1^2, m_2^2, x)} \frac{\partial}{\partial x} g^{(j)}(m_1^2, m_2^2, x) \right. \\ &\quad \left. + (m_1^2 + m_2^2 - m_3^2) g^{(j)}(m_1^2, m_2^2, m_3^2) \right\}; \end{aligned} \quad (35)$$

that last formula, as can be easily verified, satisfies the differential equations (29,30,31), with the initial conditions Eq.(32), implied by the regularity at $m_3 = m_1 + m_2$.

The integration can be actually carried out by using the change of variable

$$x = m_1^2 + m_2^2 + m_1 m_2 \frac{t^2 + 1}{t} = \frac{m_1 m_2}{t} \left(t + \frac{m_1}{m_2} \right) \left(t + \frac{m_2}{m_1} \right), \quad (36)$$

which can be inverted into

$$\begin{aligned} t = t(x; m_1^2, m_2^2) &= \frac{\sqrt{x - (m_1 - m_2)^2} + \sqrt{x - (m_1 + m_2)^2}}{\sqrt{x - (m_1 - m_2)^2} - \sqrt{x - (m_1 + m_2)^2}} \\ &= \frac{1}{2m_1 m_2} \left(x - m_1^2 - m_2^2 + R(x, m_1^2, m_2^2) \right). \end{aligned} \quad (37)$$

The solutions of the equations (29,30,31) with the initial conditions given by the equations (32) are then respectively

$$V^{(-2)}(m_1^2, m_2^2, m_3^2) = -\frac{1}{8}(m_1^2 + m_2^2 + m_3^2), \quad (38)$$

$$V^{(-1)}(m_1^2, m_2^2, m_3^2) = \frac{1}{8} \left\{ \frac{3}{2} (m_1^2 + m_2^2 + m_3^2) - \left[m_1^2 \log(m_1^2) + m_2^2 \log(m_2^2) + m_3^2 \log(m_3^2) \right] \right\}, \quad (39)$$

$$V^{(0)}(m_1^2, m_2^2, m_3^2) = \frac{1}{16} \left\{ R(m_1^2, m_2^2, m_3^2) L(m_1^2, m_2^2, m_3^2) - \frac{7}{2} (m_1^2 + m_2^2 + m_3^2) + 3 \left[m_1^2 \log(m_1^2) + m_2^2 \log(m_2^2) + m_3^2 \log(m_3^2) \right] - \frac{1}{2} \left[m_1^2 \log^2(m_1^2) + m_2^2 \log^2(m_2^2) + m_3^2 \log^2(m_3^2) \right] + \frac{1}{2} \left[(-m_1^2 - m_2^2 + m_3^2) \log(m_1^2) \log(m_2^2) + (-m_1^2 + m_2^2 - m_3^2) \log(m_1^2) \log(m_3^2) + (m_1^2 - m_2^2 - m_3^2) \log(m_2^2) \log(m_3^2) \right] \right\}, \quad (40)$$

where

$$L(m_1^2, m_2^2, m_3^2) = \text{Li}_2(-t_3 m_2/m_1) + \text{Li}_2(-t_3 m_1/m_2) + \zeta(2) + \frac{1}{2} \log^2(t_3) + \frac{1}{2} \left[\log(t_3 + m_2/m_1) - \log(t_3 + m_1/m_2) + \frac{3}{4} \log(m_1^2/m_2^2) \right] \log(m_1^2/m_2^2), \quad (41)$$

with $t_3 = t(m_3^2; m_1^2, m_2^2)$, see Eq.(37). It is to be noted that Eq.s(38-40) are expected to be symmetrical in the exchange of m_1^2, m_2^2, m_3^2 ; the symmetry, which is manifest for most terms, holds also for $L(m_1^2, m_2^2, m_3^2)$, Eq.(41), even if in this case the symmetry is not manifest. Note also that Eq.(41) remains valid in its present form for any real positive value of m_1, m_2 and m_3 , provided that t_3 is given by the second of Eq.s(37). In the limiting case $m_1, m_2 \rightarrow 0$, finally, independently of the order of the limits one finds

$$V^{(0)}(0, 0, m^2) = -\frac{m^2}{16} \left[\zeta(2) + \frac{7}{2} - 3 \log(m^2) + \log^2(m^2) \right]. \quad (42)$$

Analytical values of $V(n, m_1^2, m_2^2, m_3^2)$ were already presented in [5], where a differential equations method based on mass derivatives [6] was used, and in [7], where the results were obtained by using a Mellin-Barnes representation and its subsequent expansion in $n \rightarrow 4$ limit. Our results agree numerically with those presented in [5] in terms of Lobachevskiy's function and a related one; to check that it agrees also with the analytic formula of [7] it is necessary to use the following relation between dilogarithms of different arguments

$$\text{Li}_2(-t_2 m_1/m_3) + \text{Li}_2(-t_1 m_2/m_3) - \log(-t_2 m_1/m_3) \log(-t_1 m_2/m_3) =$$

$$\begin{aligned} & \text{Li}_2(-t_3 m_1/m_2) + \text{Li}_2(-t_3 m_2/m_1) + 2\zeta(2) + \frac{1}{2} \log^2(t_3) - \frac{1}{2} \log(m_1^2/m_3^2) \log(m_2^2/m_3^2) \\ & + \frac{1}{2} [\log(t_3 + m_2/m_1) - \log(t_3 + m_1/m_2)] \log(m_1^2/m_2^2) + \frac{3}{8} \log^2(m_1^2/m_2^2) , \end{aligned} \quad (43)$$

valid for $m_3 \geq m_1 + m_2$, with $t_1 = t(m_1^2; m_2^2, m_3^2)$ and $t_2 = t(m_2^2; m_1^2, m_3^2)$, whose validity can be established by standard methods.

4 The expansion at $p^2 \simeq 0$.

As $p^2 = 0$ is a regular point for the 4 functions $F_i(n, m_1^2, m_2^2, m_3^2, p^2)$, $i = 0, 1, 2, 3$, they can be expanded in powers of p^2 around that point

$$F_i(n, m_1^2, m_2^2, m_3^2, p^2) = \sum_{k=0}^{\infty} F_{i,k}(n, m_1^2, m_2^2, m_3^2) (p^2)^k . \quad (44)$$

The very first term, according to Eq.(24) is

$$F_{0,0}(n, m_1^2, m_2^2, m_3^2) = F_0(n, m_1^2, m_2^2, m_3^2, 0) = V(n, m_1^2, m_2^2, m_3^2) , \quad (45)$$

where the vacuum amplitude $V(n, m_1^2, m_2^2, m_3^2)$ discussed and evaluated in the previous Section, Eq.(28) and Eq.s(38-40). Once $V(n, m_1^2, m_2^2, m_3^2)$ is known, Eq.(25) and Eq.(26) can be combined to give, if (i, j, k) is any permutation of the indices $(1, 2, 3)$

$$\begin{aligned} F_{i,0}(n, m_1^2, m_2^2, m_3^2) &= F_i(n, m_1^2, m_2^2, m_3^2, 0) \\ &= -\frac{1}{R^2(m_1^2, m_2^2, m_3^2)} \left[(n-3)(m_i^2 - m_j^2 - m_k^2)V(n, m_1^2, m_2^2, m_3^2) \right. \\ &\quad + (n-2) \frac{m_i^2 - m_j^2 + m_k^2}{2m_i^2} T(n, m_j^2) T(n, m_i^2) \\ &\quad + (n-2) \frac{m_i^2 + m_j^2 - m_k^2}{2m_i^2} T(n, m_k^2) T(n, m_i^2) \\ &\quad \left. - (n-2) T(n, m_j^2) T(n, m_k^2) \right] , \end{aligned} \quad (46)$$

showing that the three quantities $F_{i,0}(n, m_1^2, m_2^2, m_3^2)$, $i = 1, 2, 3$, are all expressed algebraically in terms of $V(n, m_1^2, m_2^2, m_3^2)$ only.

When the expansion Eq.(44) is inserted in the Eq.s (4) and (7), by imposing the equality of the coefficients of the various powers of p^2 one obtains a set of equations for the quantities $F_{i,k}(n, m_1^2, m_2^2, m_3^2)$. For $k = 0$ one recovers of course the equations discussed and

solved in the previous Section; for $k > 0$ one gets a set of linear algebraic equations for the $F_{i,k}(n, m_1^2, m_2^2, m_3^2)$, which can be solved recursively in terms of the $F_{i,k-1}(n, m_1^2, m_2^2, m_3^2)$. Therefore everything can be expressed in terms of the $F_{i,0}(n, m_1^2, m_2^2, m_3^2)$, which in turn depend, as already remarked, on the single vacuum amplitude $V(n, m_1^2, m_2^2, m_3^2)$; as a consequence, all the $F_{i,k}(n, m_1^2, m_2^2, m_3^2)$ can be expressed in terms of $V(n, m_1^2, m_2^2, m_3^2)$ only.

The first iteration of the just sketched procedure gives

$$\begin{aligned}
F_{0,1}(n, m_1^2, m_2^2, m_3^2) = & \frac{n-3}{nR^4(m_1^2, m_2^2, m_3^2)} \left[(n-4) \left(m_1^4(m_2^2 + m_3^2 - m_1^2) \right. \right. \\
& + m_2^4(m_1^2 - m_2^2 + m_3^2) + m_3^4(m_1^2 + m_2^2 - m_3^2) \left. \right) - 2(n-8)m_1^2m_2^2m_3^2 \Big] V(n, m_1^2, m_2^2, m_3^2) \\
& + \frac{n-2}{nR^4(m_1^2, m_2^2, m_3^2)} \left[(n-4)(m_3^2 + m_1^2 - m_2^2)(m_3^2 - m_1^2 + m_2^2) \right. \\
& \quad \left. + 2m_3^2(m_1^2 + m_2^2 - m_3^2) \right] T(n, m_1^2)T(n, m_2^2) \\
& + \frac{n-2}{nR^4(m_1^2, m_2^2, m_3^2)} \left[(n-4)(m_2^2 + m_1^2 - m_3^2)(m_2^2 - m_1^2 + m_3^2) \right. \\
& \quad \left. + 2m_2^2(m_1^2 - m_2^2 + m_3^2) \right] T(n, m_1^2)T(n, m_3^2) \\
& + \frac{n-2}{nR^4(m_1^2, m_2^2, m_3^2)} \left[(n-4)(m_1^2 + m_2^2 - m_3^2)(m_1^2 - m_2^2 + m_3^2) \right. \\
& \quad \left. + 2m_1^2(m_2^2 - m_1^2 + m_3^2) \right] T(n, m_2^2)T(n, m_3^2) . \quad (47)
\end{aligned}$$

As already done for $V(n, m_1^2, m_2^2, m_3^2)$ in Eq.(28), the $F_{i,k}(n, m_1^2, m_2^2, m_3^2)$ can be in turn expanded in $(n-4)$ as

$$\begin{aligned}
F_{i,k}(n, m_1^2, m_2^2, m_3^2) = & C^2(n) \left\{ \frac{1}{(n-4)^2} F_{i,k}^{(-2)}(m_1^2, m_2^2, m_3^2) + \frac{1}{(n-4)} F_{i,k}^{(-1)}(m_1^2, m_2^2, m_3^2) \right. \\
& \left. + F_{i,k}^{(0)}(m_1^2, m_2^2, m_3^2) + \mathcal{O}(n-4) \right\} , \quad (48)
\end{aligned}$$

where $C(n)$ is given as usual by Eq.(9) and Eq.(11). From Eq.(47) and Eq.s(38-40) one obtains

$$F_{0,1}^{(-2)}(m_1^2, m_2^2, m_3^2) = 0 , \quad (49)$$

$$F_{0,1}^{(-1)}(m_1^2, m_2^2, m_3^2) = \frac{1}{32} , \quad (50)$$

$$F_{0,1}^{(0)}(m_1^2, m_2^2, m_3^2) = \frac{1}{32} \left\{ \frac{4m_1^2m_2^2m_3^2}{R^3(m_1^2, m_2^2, m_3^2)} L(m_1^2, m_2^2, m_3^2) - \frac{3}{4} \right.$$

$$\begin{aligned}
& + \frac{1}{R^2(m_1^2, m_2^2, m_3^2)} \left[m_1^2(m_1^2 - m_2^2 - m_3^2) \log(m_1^2) \right. \\
& \left. + m_2^2(-m_1^2 + m_2^2 - m_3^2) \log(m_2^2) + m_3^2(-m_1^2 - m_2^2 + m_3^2) \log(m_3^2) \right] \Big\} . \quad (51)
\end{aligned}$$

This result is in numerical agreement with the first term in the p^2/m_3^2 expansion in [8]. It is straightforward to obtain, when required, any other term of the double expansion in p^2 and $(n-4)$.

5 The expansion in $(n-4)$ for arbitrary p^2 .

Quite in general, by using the same argument leading to the expansion of Eq.(28), the functions $F_i(n, m_1^2, m_2^2, m_3^2, p^2)$, $i = 0, 1, 2, 3$, are expected to develop up to a double pole in $(n-4)$ and can be expanded, also for arbitrary p^2 , in the by now customary form

$$\begin{aligned}
F_i(n, m_1^2, m_2^2, m_3^2, p^2) = C^2(n) \Big\{ & \frac{1}{(n-4)^2} F_i^{(-2)}(m_1^2, m_2^2, m_3^2, p^2) \\
& + \frac{1}{(n-4)} F_i^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) + F_i^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + \mathcal{O}(n-4) \Big\} \quad (52)
\end{aligned}$$

When all the terms of Eq.s(5, 7) are consistently expanded in $(n-4)$, one obtains a set of equations for the residua and the finite part in $(n-4)$. Those equations can then be solved recursively, starting from the double pole. The equations corresponding to the poles in $(n-4)$ will not be written explicitly here for the sake of brevity; it is however easy to verify that the solutions of those equations, corresponding to the coefficients of the poles in $(n-4)$ of $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ are

$$\begin{aligned}
F_0^{(-2)}(m_1^2, m_2^2, m_3^2, p^2) &= -\frac{1}{8}(m_1^2 + m_2^2 + m_3^2) , \\
F_0^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) &= \frac{1}{8} \left\{ \frac{p^2}{4} + \frac{3}{2}(m_1^2 + m_2^2 + m_3^2) \right. \\
&\quad \left. - [m_1^2 \log(m_1^2) + m_2^2 \log(m_2^2) + m_3^2 \log(m_3^2)] \right\} , \quad (53)
\end{aligned}$$

from which the corresponding coefficients $F_i^{(k)}(m_1^2, m_2^2, m_3^2, p^2)$, for $i = 1, 2, 3$ and $k = -2, -1$ are immediately obtained by using the Eq.s(3),

$$\begin{aligned}
F_i^{(-2)}(m_1^2, m_2^2, m_3^2, p^2) &= \frac{1}{8} , \\
F_i^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) &= -\frac{1}{16} + \frac{1}{8} \log(m_i^2) . \quad (54)
\end{aligned}$$

The above functions, indeed, satisfy the differential equations and the initial condition at $p^2 = 0$ provided by Eq.s(38-40), therefore they are the exact solutions for arbitrary p^2 .

On account of the above results for the singular parts, from Eq.s(5, 7) one obtains for the finite parts $F_i^{(0)}(m_1^2, m_2^2, m_3^2, p^2)$ of the 4 amplitudes the equations

$$p^2 \frac{\partial}{\partial p^2} F_0^{(0)}(m_1^2, m_2^2, m_3^2, p^2) = F_0^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + F_0^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \\ + m_1^2 F_1^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + m_2^2 F_2^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + m_3^2 F_3^{(0)}(m_1^2, m_2^2, m_3^2, p^2) , \quad (55)$$

and

$$8D(m_1^2, m_2^2, m_3^2, p^2) p^2 \frac{\partial}{\partial p^2} F_i^{(0)}(m_1^2, m_2^2, m_3^2, p^2) = \\ 4D(m_1^2, m_2^2, m_3^2, p^2) F_i^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \\ + P_{i,0}(m_1^2, m_2^2, m_3^2, p^2) \left[16F_0^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + 28F_0^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \right. \\ \left. + 12F_0^{(-2)}(m_1^2, m_2^2, m_3^2, p^2) \right] \\ + 8P_{i,i}(m_1^2, m_2^2, m_3^2, p^2) \left[F_i^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + F_i^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \right] \\ + 8P_{i,j}(m_1^2, m_2^2, m_3^2, p^2) \left[F_j^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + F_j^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \right] \\ + 8P_{i,k}(m_1^2, m_2^2, m_3^2, p^2) \left[F_k^{(0)}(m_1^2, m_2^2, m_3^2, p^2) + F_k^{(-1)}(m_1^2, m_2^2, m_3^2, p^2) \right] \\ + Q_{i,i}(m_1^2, m_2^2, m_3^2, p^2) m_j^2 m_k^2 \left[\log(m_j^2) + \log(m_k^2) \right]^2 \\ + Q_{i,j}(m_1^2, m_2^2, m_3^2, p^2) m_i^2 m_k^2 \left[\log(m_i^2) + \log(m_k^2) \right]^2 \\ + Q_{i,k}(m_1^2, m_2^2, m_3^2, p^2) m_i^2 m_j^2 \left[\log(m_i^2) + \log(m_j^2) \right]^2 , \quad (56)$$

where (i, j, k) is any permutation of $(1, 2, 3)$ and $P_{i,j}(m_1^2, m_2^2, m_3^2, p^2)$, $Q_{i,j}(m_1^2, m_2^2, m_3^2, p^2)$ are the polynomials of Eq.(7).

In the notation of Eq.(52), for $j = -2, -1, 0$

$$F_0^{(j)}(m_1^2, m_2^2, m_3^2, 0) = V^{(j)}(m_1^2, m_2^2, m_3^2) , \quad (57)$$

where the $V^{(j)}(m_1^2, m_2^2, m_3^2)$ are given in Eq.s(38-40), so that $F_0^{(0)}(m_1^2, m_2^2, m_3^2, 0)$ is known; similarly, the $F_i^{(0)}(m_1^2, m_2^2, m_3^2, 0)$, $i = 1, 2, 3$, can be obtained by using either Eq.(25) or Eq.(46) (properly expanded in the $n \rightarrow 4$ limit). As a conclusion, Eq.(55) and Eq.(56), together with the initial conditions at $p^2 = 0$ just discussed, are free by construction from the singularities of the $n \rightarrow 4$ limit and therefore ready to be used for the numerical evaluation of the finite parts of the sunrise master integrals.

For convenience of later use, we introduce one more quantity $\mathcal{F}_0(m_1^2, m_2^2, m_3^2, p^2)$, defined by

$$\mathcal{F}_0(m_1^2, m_2^2, m_3^2, p^2) = \lim_{n \rightarrow 4} \left[F_0(n, m_1^2, m_2^2, m_3^2, p^2) - F_{0,0}(n, m_1^2, m_2^2, m_3^2) - p^2 F_{0,1}(n, m_1^2, m_2^2, m_3^2) \right], \quad (58)$$

where $F_{0,0}(n, m_1^2, m_2^2, m_3^2)$ and $F_{0,1}(n, m_1^2, m_2^2, m_3^2)$ are given by Eq.(45) and Eq.(47).

It is to be noted that the terms of $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ singular in $(n-4)$ are at most linear in p^2 , as shown in Eq.s(53), and therefore entirely encapsulated in the combination $F_{0,0}(n, m_1^2, m_2^2, m_3^2) + p^2 F_{0,1}(n, m_1^2, m_2^2, m_3^2)$; for that reason $\mathcal{F}_0(m_1^2, m_2^2, m_3^2, p^2)$ is finite in the $n \rightarrow 4$ limit.

From the very definition Eq.(58), $\mathcal{F}_0(m_1^2, m_2^2, m_3^2, p^2)$ has a double zero at $p^2 = 0$; therefore it satisfies the doubly subtracted dispersion relation

$$\mathcal{F}_0(m_1^2, m_2^2, m_3^2, p^2) = (p^2)^2 \int_{(m_1+m_2+m_3)^2}^{\infty} \frac{du}{u^2(u+p^2)} \frac{1}{\pi} \text{Im} F_0(4, m_1^2, m_2^2, m_3^2, -u), \quad (59)$$

which is obviously finite at $n = 4$.

6 The analytic calculation for $m_2 = m_3 = 0$.

We discuss now the case in which all the masses but one are equal to zero; the remarkably simpler equations obtained in that limit can be solved analytically, providing useful information for the study of the large p^2 behaviour, even for arbitrary masses, which will be worked out in the next Section.

When putting $m_2 = m_3 = 0$, and $m_1 = m$ for simplicity, the system of the 4 equations Eq.(5) and Eq.(7) collapses into just two equations:

$$\begin{aligned} p^2 \frac{\partial}{\partial p^2} F_0(n, m^2, 0, 0, p^2) &= m^2 F_1(n, m^2, 0, 0, p^2) + (n-3) F_0(n, m^2, 0, 0, p^2) \\ p^2 (p^2 + m^2) \frac{\partial}{\partial p^2} F_1(n, m^2, 0, 0, p^2) &= -\frac{1}{2} (n-3) (3n-8) F_0(n, m^2, 0, 0, p^2) \\ &\quad + \left[2p^2 (n-3) - (p^2 + m^2) \left(\frac{3}{2}n - 4 \right) \right] F_1(n, m^2, 0, 0, p^2). \end{aligned} \quad (60)$$

The system so obtained is homogeneous, and as such it cannot provide information on the initial values; but the required initial conditions are still supplied by the proper $m_2 = m_3 = 0$ limit of the results of the previous sections.

To solve the system, we observe that it is of course equivalent to a second order differential equation in a single function. If $F_0(n, m^2, 0, 0, p^2)$ is kept as the independent function, eliminating $F_1(n, m^2, 0, 0, p^2)$ from Eq.s(60) one obtains

$$p^2(p^2 + m^2) \frac{\partial^2}{\partial (p^2)^2} F_0(n, m^2, 0, 0, p^2) + \left(-2p^2(n-3) + \frac{n}{2}(p^2 + m^2) \right) \frac{\partial}{\partial p^2} F_0(n, m^2, 0, 0, p^2) + \frac{(n-4)(n-3)}{2} F_0(n, m^2, 0, 0, p^2) = 0 . \quad (61)$$

By introducing $\mathcal{F}_0(m^2, 0, 0, p^2)$ according to Eq.(58), with the values of $F_{0,0}(n, 0, 0, m^2)$ and $F_{0,1}(n, 0, 0, m^2)$ corresponding to Eq.s(45),(47) in the $m_1 = m_2 = 0$ limit given by Eq.(28) and Eq.(42), one has

$$F_0(n, m^2, 0, 0, p^2) = C^2(n) \left\{ -\frac{1}{(n-4)^2} \frac{m^2}{8} + \frac{1}{(n-4)} \frac{1}{32} (p^2 + 6m^2 - 4m^2 \log(m^2)) + \frac{1}{16} \left[p^2 \left(-\frac{3}{8} + \frac{1}{2} \log(m^2) \right) - m^2 \left(\zeta(2) + \frac{7}{2} - 3 \log(m^2) + \log^2(m^2) \right) \right] + \mathcal{F}_0(m^2, 0, 0, p^2) + \dots \right\} . \quad (62)$$

In terms of $\mathcal{F}_0(m^2, 0, 0, p^2)$ Eq.(61) then reads

$$p^2(p^2 + m^2) \frac{\partial^2}{\partial (p^2)^2} \mathcal{F}_0(m^2, 0, 0, p^2) + 2m^2 \frac{\partial}{\partial p^2} \mathcal{F}_0(m^2, 0, 0, p^2) = \frac{1}{32} p^2 , \quad (63)$$

which is in fact a first order differential equation in $\frac{\partial}{\partial p^2} \mathcal{F}_0(m^2, 0, 0, p^2)$. The values of $\mathcal{F}_0(m^2, 0, 0, p^2)$ and its derivative at $p^2 = 0$ are by construction

$$\begin{aligned} \mathcal{F}_0(m^2, 0, 0, 0) &= 0 , \\ \frac{\partial}{\partial p^2} \mathcal{F}_0(m^2, 0, 0, 0) &= 0 . \end{aligned} \quad (64)$$

With the substitution

$$\frac{\partial}{\partial p^2} \mathcal{F}_0(m^2, 0, 0, p^2) = \frac{(p^2 + m^2)^2}{(p^2)^2} G(m^2, p^2) , \quad (65)$$

Eq.(63) becomes

$$\frac{\partial}{\partial p^2} G(m^2, p^2) = \frac{1}{32} \frac{(p^2)^2}{(p^2 + m^2)^3} . \quad (66)$$

The integration of Eq.(66) is trivial; the result in terms of $\mathcal{F}_0(m^2, 0, 0, p^2)$, accounting for the second of the initial condition of Eq.(64), reads

$$\frac{\partial}{\partial p^2} \mathcal{F}_0(m^2, 0, 0, p^2) = \frac{1}{32} \left[\frac{(p^2 + m^2)^2}{(p^2)^2} \log\left(1 + \frac{p^2}{m^2}\right) - \frac{m^2}{p^2} - \frac{3}{2} \right] . \quad (67)$$

One more integration and the first initial condition of Eq.(64) give

$$\mathcal{F}_0(m^2, 0, 0, p^2) = \frac{1}{32} \left\{ -2m^2 \text{Li}_2 \left(-\frac{p^2}{m^2} \right) + \left(p^2 - \frac{m^4}{p^2} \right) \log \left(1 + \frac{p^2}{m^2} \right) - \frac{5}{2} p^2 + m^2 \right\} . \quad (68)$$

The same result can be obtained, by means of the doubly subtracted dispersion relations Eq.(59) for the specific values of the masses,

$$\mathcal{F}_0(m^2, 0, 0, p^2) = (p^2)^2 \int_{m^2}^{\infty} \frac{du}{u^2(u + p^2)} \frac{1}{\pi} \text{Im} F_0(4, m^2, 0, 0, -u) . \quad (69)$$

The Cutkosky-Veltman rule gives for the imaginary part

$$\begin{aligned} \frac{1}{\pi} \text{Im} F_0(4, m^2, 0, 0, -u) &= \frac{1}{16} \int_{m^2}^u db \frac{u-b}{u} \frac{b-m^2}{b} \\ &= \frac{1}{16} \left\{ \frac{u^2 - m^4}{2u} - m^2 \log \frac{u}{m^2} \right\} ; \end{aligned} \quad (70)$$

substituting in Eq.(69) and carrying out a last integration Eq.(68) is recovered.

7 Large p^2 expansion.

By using the differential equations (5,7) it is also possible to study the asymptotic expansion of the 4 functions $F_i(n, m_1^2, m_2^2, m_3^2, p^2)$ for large values of p^2 . As $p^2 = \infty$ is a singular point of the equations, we look for a tentative asymptotic expansion of F_0 in the form

$$F_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2, p^2) = (p^2)^\alpha \sum_{k=0}^{\infty} \Phi_k^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) \frac{1}{(p^2)^k} . \quad (71)$$

Due to the linearity of the equations, the actual solution is built by summing a suitable linear combination of all the possible expansions. The corresponding expansions of the other 3 functions $F_i^{(\alpha)}(n, m_1^2, m_2^2, m_3^2, p^2)$, $i = 1, 2, 3$, can be obtained through Eq.(3).

By inserting the expansion Eq.(71) into Eq.(5) and Eq.(7)), one obtains a set of 4 equations, which must be satisfied, for each separate power of p^2 , by the coefficients of the expansion.

A first solution corresponds to $\alpha = -1$; we write it as

$$\begin{cases} \alpha = -1 \equiv r , \\ \frac{1}{p^2} \sum_{k=0}^{\infty} \phi_k^{(r)}(n, m_1^2, m_2^2, m_3^2) \frac{1}{(p^2)^k} . \end{cases} \quad (72)$$

It corresponds to the so-called regular solution, hence the superscript (r) in $\phi_k^{(r)}$ (we write $\phi_k^{(r)}$ rather than $\Phi_k^{(r)}$ for uniformity with later use). Its coefficients are completely determined by the non-homogeneous part of the equations; correspondingly, all the terms behave for large p^2 as strictly integer powers of $1/p^2$, which is exactly the behaviour of the various terms occurring in the asymptotic expansion of the non-homogeneous part. The first coefficient of the expansion is

$$\phi_0^{(r)}(n, m_1^2, m_2^2, m_3^2) = T(n, m_1^2)T(n, m_2^2) + T(n, m_1^2)T(n, m_3^2) + T(n, m_2^2)T(n, m_3^2) ; \quad (73)$$

all the other coefficients $\phi_k^{(r)}$ can be similarly obtained, when needed, by following the procedure just outlined.

Other solutions correspond to different values of α ; if α is not equal to -1 , the leading coefficient must satisfy the homogeneous system of equations

$$\begin{aligned} \sum_{i=1}^3 m_i^2 \frac{\partial}{\partial m_i^2} \Phi_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) + (\alpha - n + 3) \Phi_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) &= 0 \\ \left[\alpha - \frac{1}{2}(n - 4) \right] \frac{\partial}{\partial m_1^2} \Phi_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) &= 0 \\ \left[\alpha - \frac{1}{2}(n - 4) \right] \frac{\partial}{\partial m_2^2} \Phi_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) &= 0 \\ \left[\alpha - \frac{1}{2}(n - 4) \right] \frac{\partial}{\partial m_3^2} \Phi_0^{(\alpha)}(n, m_1^2, m_2^2, m_3^2) &= 0 . \end{aligned} \quad (74)$$

There are two solutions of the system (74), corresponding to the values $\alpha = (n - 3)$ and $\alpha = (n - 4)/2$. Correspondingly, Eq.s(74) become

$$\begin{cases} \alpha = n - 3 \equiv s_1 , \\ \frac{\partial}{\partial m_i^2} \Phi_0^{(s_1)}(n, m_1^2, m_2^2, m_3^2) = 0 \quad (i = 1, 2, 3) , \end{cases} \quad (75)$$

and

$$\begin{cases} \alpha = \frac{1}{2}(n - 4) \equiv s_2 , \\ \sum_{i=1}^3 m_i^2 \frac{\partial}{\partial m_i^2} \Phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) - \frac{n-2}{2} \Phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) = 0 ; \end{cases} \quad (76)$$

the above values of α are also called “fractionary”, and the corresponding solutions singular, hence the superscripts $(s_1), (s_2)$, as opposed to the superscript (r) for the “regular” solution corresponding to the integer value $\alpha = -1$ of Eq.(72). Due to the homogeneity of the equations for the coefficients of the “fractional” powers, the two leading coefficients $\Phi_0^{(s_i)}(n, m_1^2, m_2^2, m_3^2)$ cannot be determined by the equations (75) and (76) alone; nevertheless, the equations can still provide much valuable information.

Eq.(75) shows indeed that $\Phi_0^{(s_1)}(n, m_1^2, m_2^2, m_3^2)$ is independent of the masses

$$\Phi_0^{(s_1)}(n, m_1^2, m_2^2, m_3^2) = \Phi^{(s_1)}(n) , \quad (77)$$

so that the still unknown function $\Phi^{(s_1)}(n)$ depends only on n .

The case of Eq.(76) is slightly more complicated; by investigating the condition on the leading $(p^2)^{(n-4)/2}$ term implied by Eq.(95) of the Appendix, whose *l.h.s.* is the second derivative of $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ with respect to m_1^2 and m_2^2 , one finds the relations

$$\frac{\partial^2}{\partial m_i^2 \partial m_j^2} \Phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) = 0 \quad (i, j = 1, 2, 3; i \neq j) . \quad (78)$$

According to Eq.(78), the solution of Eq.(76) can be written as

$$\begin{aligned} \Phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) &= \Phi^{(s_2)}(n) \phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) , \\ \phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) &= m_1^{n-2} + m_2^{n-2} + m_3^{n-2} , \end{aligned} \quad (79)$$

where $\Phi^{(s_2)}(n)$ is a function of n only.

For each “fractional” power solution, all the next-to-leading terms of the expansion are determined in terms of the leading terms, by solving recursively the systems of linear equations; it is therefore convenient to factorize the (as yet arbitrary) function $\Phi^{(s_i)}(n)$, defining

$$\Phi_k^{(s_i)}(n, m_1^2, m_2^2, m_3^2) = \Phi^{(s_i)}(n) \phi_k^{(s_i)}(n, m_1^2, m_2^2, m_3^2) . \quad (80)$$

An explicit calculation then gives for the first few terms (which we will use later)

$$\begin{aligned} \phi_1^{(s_1)}(n, m_1^2, m_2^2, m_3^2) &= \frac{(n-3)(3n-8)}{(n-4)} (m_1^2 + m_2^2 + m_3^2) , \\ \phi_2^{(s_1)}(n, m_1^2, m_2^2, m_3^2) &= \frac{(n-3)(3n-8)}{(n-6)} \left[-\frac{1}{(n-4)} B(m_1^2, m_2^2, m_3^2) + R^2(m_1^2, m_2^2, m_3^2) \right. \\ &\quad \left. - \frac{1}{2} B(m_1^2, m_2^2, m_3^2) + \frac{3}{2} (n-4) \left(R^2(m_1^2, m_2^2, m_3^2) + \frac{1}{2} B(m_1^2, m_2^2, m_3^2) \right) \right] , \end{aligned} \quad (81)$$

where

$$B(m_1^2, m_2^2, m_3^2) = 4(m_1^2 m_2^2 + m_1^2 m_3^2 + m_2^2 m_3^2) , \quad (82)$$

and

$$\begin{aligned} \phi_1^{(s_2)}(n, m_1^2, m_2^2, m_3^2) = & \frac{n-3}{n} \left[-(3n-8)(m_1^2 + m_2^2 + m_3^2) \phi_0^{(s_2)}(n, m_1^2, m_2^2, m_3^2) \right. \\ & + 2(n-2) \left(m_1^{n-2}(m_1^2 + 2m_2^2 + 2m_3^2) + m_2^{n-2}(2m_1^2 + m_2^2 + 2m_3^2) \right. \\ & \left. \left. + m_3^{n-2}(2m_1^2 + 2m_2^2 + m_3^2) \right) \right] . \end{aligned} \quad (83)$$

Any subsequent term can be easily obtained, when needed, by the same procedure.

As for arbitrary n the three solutions satisfy the equations independently of each other, the actual asymptotic expansion of $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ can be written as their sum

$$\begin{aligned} F_0^{(\infty)}(n, m_1^2, m_2^2, m_3^2, p^2) = & p^2 (p^2)^{n-4} \Phi^{(s_1)}(n) \left(1 + \sum_{k=1}^{\infty} \phi_k^{(s_1)}(n, m_1^2, m_2^2, m_3^2) \frac{1}{(p^2)^k} \right) \\ & + (p^2)^{\frac{1}{2}(n-4)} \Phi^{(s_2)}(n) \sum_{k=0}^{\infty} \phi_k^{(s_2)}(n, m_1^2, m_2^2, m_3^2) \frac{1}{(p^2)^k} \\ & + \frac{1}{p^2} \sum_{k=0}^{\infty} \phi_k^{(r)}(n, m_1^2, m_2^2, m_3^2) \frac{1}{(p^2)^k} ; \end{aligned} \quad (84)$$

the above expression depends on the two functions of n , $\Phi^{(s_1)}(n)$ and $\Phi^{(s_2)}(n)$, which are as yet undetermined. To fix them, we must supply more information on $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$. As the $F_i(n, m_1^2, m_2^2, m_3^2, p^2)$, $i = 0, 1, 2, 3$, were studied in the $n \rightarrow 4$ limit, Eq.(52), we expand also Eq.(84) around $n = 4$, writing in particular $\Phi^{(s_1)}(n)$ and $\Phi^{(s_2)}(n)$ as

$$\Phi^{(s_i)}(n) = C^2(n) \sum_{l=-2}^{\infty} (n-4)^l \Phi^{(s_i, l)} \quad (i = 1, 2) ; \quad (85)$$

as always $C(n)$ is not expanded, appearing as an overall factor.

By comparison with the coefficients of the singularities in $(n-4)$, which are exactly known and given in Eq.(53), the following values are found

$$\begin{aligned} \Phi^{(s_1, -2)} &= 0 , & \Phi^{(s_2, -2)} &= -\frac{1}{4} \\ \Phi^{(s_1, -1)} &= \frac{1}{32} , & \Phi^{(s_2, -1)} &= -4\Phi^{(s_1, 0)} - \frac{1}{32} . \end{aligned} \quad (86)$$

Unfortunately some additional information has still to be supplied, besides that coming from the singularities in $(n-4)$, for fixing the other as yet unknown constants $\Phi^{(s_1, 0)}$,

$\Phi^{(s_2,0)}$ and $\Phi^{(s_1,1)}$; the last quantity is the coefficient of the term in $(n-4)$ in Eq.(85), which is also needed due to the presence of the $1/(n-4)$ factor in Eq.(81).

As those constants are independent of the masses, they can be determined through the comparison with the asymptotic expansion of a simpler case, namely $F_0(n, m^2, 0, 0, p^2)$, discussed in the previous Section, whose explicit analytic value is given in Eq.(68). Its large p^2 expansion reads

$$\begin{aligned} F_0^{(\infty)}(n, m^2, 0, 0, p^2) = & C^2(n) \left\{ -\frac{1}{(n-4)^2} \frac{m^2}{8} + \frac{1}{(n-4)} \frac{1}{32} (p^2 + 6m^2 - 4m^2 \log(m^2)) \right. \\ & + \frac{1}{16} \left[p^2 \left(-\frac{13}{8} + \frac{1}{2} \log(p^2) \right) + \frac{m^2}{2} \log^2(p^2) - m^2 \log(m^2) \log(p^2) \right. \\ & + m^2 \left(-\frac{1}{2} \log^2(m^2) + 3 \log(m^2) - \frac{5}{2} \right) \\ & \left. \left. - \frac{m^4}{2p^2} \left(\log(p^2) - \log(m^2) + \frac{5}{2} \right) \right] + \dots \right\} . \end{aligned} \quad (87)$$

Let us recall that, when using Eq.(84) and Eq.(85), the logarithmic factors are generated in the $n \rightarrow 4$ limit through expansions of the kind

$$\frac{1}{(n-4)^2} (p^2)^{n-4} = \frac{1}{(n-4)^2} \left(1 + (n-4) \log(p^2) + \frac{1}{2} (n-4)^2 \log^2(p^2) + \dots \right) .$$

By comparing Eq.(87) and the $n \rightarrow 4$ limit of Eq.(84), when two masses are taken equal to zero, Eq.(86) is completed into

$$\begin{aligned} \Phi^{(s_1,-2)} &= 0 , & \Phi^{(s_2,-2)} &= -\frac{1}{4} \\ \Phi^{(s_1,-1)} &= \frac{1}{32} , & \Phi^{(s_2,-1)} &= \frac{3}{8} \\ \Phi^{(s_1,0)} &= -\frac{13}{128} , & \Phi^{(s_2,0)} + 4\Phi^{(s_1,1)} &= \frac{59}{128} . \end{aligned} \quad (88)$$

It is apparent from the above equations that only a particular combination of $\Phi^{(s_2,0)}$ and $\Phi^{(s_1,1)}$ is determined by the comparison, but only that combination is needed for expressing $F_0(n, m_1^2, m_2^2, m_3^2, p^2)$ up to the constant term in $(n-4)$ (*i.e.* the separate knowledge of the two terms is not needed up to the degree of expansion in $(n-4)$ and p^2 , considered here).

Summing up, for arbitrary masses and in the $n \rightarrow 4$ limit, up to terms $1/p^2$ included, the asymptotic expansion Eq.(84) can be written as

$$F_0^{(\infty)}(m_1^2, m_2^2, m_3^2, p^2) = C^2(n) \left\{ \frac{1}{(n-4)^2} F_0^{(\infty,-2)}(m_1^2, m_2^2, m_3^2, p^2) \right.$$

$$\begin{aligned}
& + \frac{1}{(n-4)} F_0^{(\infty, -1)}(m_1^2, m_2^2, m_3^2, p^2) \\
& + F_0^{(\infty, 0)}(m_1^2, m_2^2, m_3^2, p^2) + \dots \Big\} , \tag{89}
\end{aligned}$$

with

$$F_0^{(\infty, -2)}(m_1^2, m_2^2, m_3^2, p^2) = -\frac{1}{8} (m_1^2 + m_2^2 + m_3^2) , \tag{90}$$

$$F_0^{(\infty, -1)}(m_1^2, m_2^2, m_3^2, p^2) = \frac{1}{16} \left[\frac{1}{2} p^2 + \sum_{i=1}^3 m_i^2 (3 - 2 \log(m_i^2)) \right] , \tag{91}$$

and

$$\begin{aligned}
F_0^{(\infty, 0)}(m_1^2, m_2^2, m_3^2, p^2) = & p^2 [f_{1,1} \log(p^2) + f_{1,0}] \\
& + f_{0,2} \log^2(p^2) + f_{0,1} \log(p^2) + f_{0,0} \\
& + \frac{1}{p^2} [f_{-1,2} \log^2(p^2) + f_{-1,1} \log(p^2) + f_{-1,0}] + \dots , \tag{92}
\end{aligned}$$

where

$$\begin{aligned}
f_{1,1} &= \frac{1}{32} , \quad f_{1,0} = -\frac{13}{128} , \quad f_{0,2} = \frac{1}{32} \sum_{i=1}^3 m_i^2 , \quad f_{0,1} = -\frac{1}{16} \sum_{i=1}^3 m_i^2 \log(m_i^2) , \\
f_{0,0} &= -\frac{1}{32} \sum_{i=1}^3 m_i^2 (5 - 6 \log(m_i^2) + \log^2(m_i^2)) , \quad f_{-1,2} = \frac{1}{64} B(m_1^2, m_2^2, m_3^2) \\
f_{-1,1} &= \frac{1}{64} (B(m_1^2, m_2^2, m_3^2) - 2R^2(m_1^2, m_2^2, m_3^2) - 4(m_2^2 + m_3^2)m_1^2 \log(m_1^2) \\
& \quad - 4(m_1^2 + m_3^2)m_2^2 \log(m_2^2) - 4(m_1^2 + m_2^2)m_3^2 \log(m_3^2)) \\
f_{-1,0} &= \frac{1}{128} (-3B(m_1^2, m_2^2, m_3^2) - 10R^2(m_1^2, m_2^2, m_3^2) + 4(m_1^2 - 2m_2^2 - 2m_3^2)m_1^2 \log(m_1^2) \\
& \quad + 4(m_2^2 - 2m_1^2 - 2m_3^2)m_2^2 \log(m_2^2) + 4(m_3^2 - 2m_1^2 - 2m_2^2)m_3^2 \log(m_3^2) \\
& \quad + 8m_1^2 m_2^2 \log(m_1^2) \log(m_2^2) + 8m_1^2 m_3^2 \log(m_1^2) \log(m_3^2) \\
& \quad + 8m_2^2 m_3^2 \log(m_2^2) \log(m_3^2)) . \tag{93}
\end{aligned}$$

The coefficients of $1/(n-4)^2$ and $1/(n-4)$, Eq.(90) and Eq.(91), are of course identical to the exact ones in Eq.(53)), as their expansions in $1/p^2$ contain only a linear and a constant term. The results given in Eq.s(92,93) agree with the first terms of the corresponding expansion given in [8].

Higher order terms, when needed, can be added with little effort by following the above discussed procedure.

Acknowledgments.

One of us (HC) is grateful to the Bologna Section of INFN and to the Department of Physics of Bologna University for support and kind hospitality.

Appendix.

We list here a few ready-to-use formulae, obtained by solving the integration by part identities for the amplitudes of the sunrise self-mass graph (see also [3]). The notation for the amplitudes is given in Eq.(1). Further relations can be found by permutation of masses.

$$\begin{aligned}
& D(m_1^2, m_2^2, m_3^2, p^2) A(n, m_1^2, m_2^2, m_3^2, p^2, -3, -1, -1, 0, 0) = \\
& \quad - \frac{n-4}{4m_1^2} D(m_1^2, m_2^2, m_3^2, p^2) F_1(n, m_1^2, m_2^2, m_3^2, p^2) \\
& \quad + (n-3)(p^2 + m_1^2) \left[4(m_1^2 m_2^2 + m_1^2 m_3^2 - m_2^2 m_3^2) \right. \\
& \quad \quad \left. - (p^2 + m_1^2 + m_2^2 + m_3^2)^2 \right] F_1(n, m_1^2, m_2^2, m_3^2, p^2) \\
& + (n-3)(p^2 + m_2^2) \frac{m_2^2}{m_1^2} \left[4(m_1^2 m_2^2 - m_1^2 m_3^2 + m_2^2 m_3^2) \right. \\
& \quad \quad \left. - (p^2 + m_1^2 + m_2^2 + m_3^2)^2 \right] F_2(n, m_1^2, m_2^2, m_3^2, p^2) \\
& + (n-3)(p^2 + m_3^2) \frac{m_3^2}{m_1^2} \left[4(-m_1^2 m_2^2 + m_1^2 m_3^2 + m_2^2 m_3^2) \right. \\
& \quad \quad \left. - (p^2 + m_1^2 + m_2^2 + m_3^2)^2 \right] F_3(n, m_1^2, m_2^2, m_3^2, p^2) \\
& + (n-3)(3n-8) \frac{1}{m_1^2} \left[-4m_1^2 m_2^2 m_3^2 + (m_1^2 m_2^2 + m_1^2 m_3^2 + m_2^2 m_3^2)(p^2 + m_1^2 + m_2^2 + m_3^2) \right. \\
& \quad \quad \left. - \frac{1}{4}(p^2 + m_1^2 + m_2^2 + m_3^2)^3 \right] F_0(n, m_1^2, m_2^2, m_3^2, p^2) \\
& + \frac{(n-2)^2}{m_1^2} \left[(m_3^2 - m_1^2)(m_3^2 - m_2^2) - \frac{1}{4}(p^2 + m_1^2 + m_2^2 - m_3^2)^2 \right] T(n, m_1^2) T(n, m_2^2) \\
& + \frac{(n-2)^2}{m_1^2} \left[(m_2^2 - m_1^2)(m_2^2 - m_3^2) - \frac{1}{4}(p^2 + m_1^2 - m_2^2 + m_3^2)^2 \right] T(n, m_1^2) T(n, m_3^2) \\
& + \frac{(n-2)^2}{m_1^2} \left[(m_1^2 - m_2^2)(m_1^2 - m_3^2) - \frac{1}{4}(p^2 - m_1^2 + m_2^2 + m_3^2)^2 \right] T(n, m_2^2) T(n, m_3^2) \quad , \quad (94)
\end{aligned}$$

$$\begin{aligned}
& D(m_1^2, m_2^2, m_3^2, p^2) A(n, m_1^2, m_2^2, m_3^2, p^2, -1, -2, -2, 0, 0) = \\
& \quad 8(n-3)m_1^2(p^2 + m_1^2)(p^2 - m_1^2 + m_2^2 + m_3^2)F_1(n, m_1^2, m_2^2, m_3^2, p^2) \\
& - (n-3) \left[(p^2 - m_1^2 + m_2^2 + m_3^2)^3 - 4(m_2^2 - m_1^2)(p^2 - m_1^2 + m_2^2 + m_3^2)^2 \right. \\
& \quad \left. - 4(m_3^2 - m_1^2)(m_1^2 + m_2^2)(p^2 - m_1^2 + m_2^2 + m_3^2) \right. \\
& \quad \left. + 16m_2^2(m_3^2 - m_1^2)(m_2^2 - m_1^2) \right] F_2(n, m_1^2, m_2^2, m_3^2, p^2) \\
& - (n-3) \left[(p^2 - m_1^2 + m_2^2 + m_3^2)^3 - 4(m_3^2 - m_1^2)(p^2 - m_1^2 + m_2^2 + m_3^2)^2 \right. \\
& \quad \left. - 4(m_2^2 - m_1^2)(m_1^2 + m_3^2)(p^2 - m_1^2 + m_2^2 + m_3^2) \right. \\
& \quad \left. + 16m_3^2(m_3^2 - m_1^2)(m_2^2 - m_1^2) \right] F_3(n, m_1^2, m_2^2, m_3^2, p^2) \\
& + (n-3)(3n-8) \left[(p^2 - m_1^2 + m_2^2 + m_3^2)^2 - 4(m_3^2 - m_1^2)(m_2^2 - m_1^2) \right] F_0(n, m_1^2, m_2^2, m_3^2, p^2) \\
& - \frac{(n-2)^2}{2m_2^2} \left[(p^2 + m_1^2 - m_2^2 + m_3^2)^2 + 4(m_3^2 - m_2^2)(m_2^2 - m_1^2) \right] T(n, m_1^2) T(n, m_2^2) \\
& - \frac{(n-2)^2}{2m_3^2} \left[(p^2 + m_1^2 + m_2^2 - m_3^2)^2 + 4(m_2^2 - m_3^2)(m_3^2 - m_1^2) \right] T(n, m_1^2) T(n, m_3^2) \\
& + \frac{(n-2)^2}{4m_2^2 m_3^2} \left[(p^2 + m_1^2 + m_2^2 + m_3^2)^3 - 4(m_1^2 m_2^2 + m_2^2 m_3^2 + m_1^2 m_3^2)(p^2 + m_1^2 + m_2^2 + m_3^2) \right. \\
& \quad \left. + 16m_1^2 m_2^2 m_3^2 \right] T(n, m_2^2) T(n, m_3^2) \quad . \quad (95)
\end{aligned}$$

References

- [1] E. Remiddi, *Nuovo Cimento* **A110**, 1435 (1997), hep-th/9711188 (1997). Similar equations, for the one-loop box amplitudes in 4 dimensions, have been proposed by V. de Alfaro, B. Jakšić and T. Regge in their paper *Differential Properties of Feynman Amplitudes* to the volume *High-Energy Physics and Elementary Particles*, IAEA, Vienna 1965.
- [2] K.G. Chetyrkin and F.V. Tkachov, *Nuc. Phys.* **B192**, 159 (1981); F.V. Tkachov, *Phys. Lett.* **B100**, 65 (1981).
- [3] O.V. Tarasov, *Nuc. Phys.* **B502**, 455 (1997).
- [4] J.A.M. Vermaseren, *Symbolic Manipulation with FORM*, Computer Algebra Nederland, Amsterdam (1991).
- [5] C. Ford, I. Jack and D.R.T. Jones, *Nuc. Phys.* **B387**, 373 (1992).
- [6] A.V. Kotikov, *Phys. Lett.* **B254**, 158 (1991).
- [7] A.I. Davydychev and J.B. Tausk, *Nuc. Phys.* **B397**, 123 (1993); A.I. Davydychev, V.A. Smirnov and J.B. Tausk, *Nuc. Phys.* **B410**, 325 (1993).
- [8] F.A. Berends, M. Böhm, M. Buza and R. Scharf, *Z. Phys.* **C 63**, 227 (1994), see in particular for small p^2 Eq.s(28,31) and for large p^2 Eq.(32) therein.